

- Campos LW, Chakrabarty S, Haque R, Martin JH. Regenerating motor bridge axons refine connections and synapse on lumbar motoneurons to bypass chronic spinal cord injury. *J Comp Neurol*. 2008 Feb 10;506(5):838-50.
- Campos L, Meng Z, Hu G, Chiu DT, Ambron RT, Martin JH. Engineering novel spinal circuits to promote recovery after spinal injury. *J Neurosci*. 2004 Mar 3;24(9):2090-101.
- Conta AC, Stelzner DJ. Immunomodulatory effect of the purine nucleoside inosine following spinal cord contusion injury in rat. *Spinal Cord*. 2008 Jan;46(1):39-44.
- Conta AC, Stelzner DJ. Differential vulnerability of propriospinal tract neurons to spinal cord contusion injury. *J Comp Neurol*. 2004 Nov 22;479(4):347-59.
- Galtrey CM, Kwok JC, Carulli D, Rhodes KE, Fawcett JW. Distribution and synthesis of extracellular matrix proteoglycans, hyaluronan, link proteins and tenascin-R in the rat spinal cord. *Eur J Neurosci*. 2008 Mar;27(6):1373-90.
- Galtrey CM, Asher RA, Nothias F, Fawcett JW. Promoting plasticity in the spinal cord with chondroitinase improves functional recovery after peripheral nerve repair. *Brain*. 2007 Apr;130(Pt 4):926-39.
- Deepa SS, Carulli D, Galtrey C, Rhodes K, Fukuda J, Mikami T, Sugahara K, Fawcett JW. Composition of perineuronal net extracellular matrix in rat brain: a different disaccharide composition for the net-associated proteoglycans. *J Biol Chem*. 2006 Jun 30;281(26):17789-800.
- Okamura RM, Lebkowski J, Au M, Priest CA, Denham J, Majumdar AS. Immunological properties of human embryonic stem cell-derived oligodendrocyte progenitor cells. *J Neuroimmunol*. 2007 Dec;192(1-2):134-44.
- LaPlaca MC, Prado GR, Cullen D, Simon CM. Plasma membrane damage as a marker of neuronal injury. *Conf Proc IEEE Eng Med Biol Soc*. 2009;2009:1113-6.
- Simon CM, Sharif S, Tan RP, LaPlaca MC. Spinal cord contusion causes acute plasma membrane damage. *J Neurotrauma*. 2009 Apr;26(4):563-74
- LaPlaca MC, Simon CM, Prado GR, Cullen DK. CNS injury biomechanics and experimental models. *Prog Brain Res*. 2007;161:13-26.
- Kuerzi J, Brown EH, Shum-Siu A, Siu A, Burke D, Morehouse J, Smith RR, Magnuson DS. Task-specificity vs. ceiling effect: step-training in shallow water after spinal cord injury. *Exp Neurol*. 2010 Jul;224(1):178-87.
- Smith RR, Brown EH, Shum-Siu A, Whelan A, Burke DA, Benton RL, Magnuson DS. Swim training initiated acutely after spinal cord injury is ineffective and induces extravasation in and around the epicenter. *J Neurotrauma*. 2009 Jul;26(7):1017-27

- Magnuson DS, Smith RR, Brown EH, Enzmann G, Angeli C, Quesada PM, Burke D. Swimming as a model of task-specific locomotor retraining after spinal cord injury in the rat. *Neurorehabil Neural Repair*. 2009 Jul-Aug;23(6):535-45.
- Zhang YP, Burke DA, Shields LB, Chekmenev SY, Dincman T, Zhang Y, Zheng Y, Smith RR, Benton RL, DeVries WH, Hu X, Magnuson DS, Whittemore SR, Shields CB. Spinal cord contusion based on precise vertebral stabilization and tissue displacement measured by combined assessment to discriminate small functional differences. *J Neurotrauma*. 2008 Oct;25(10):1227-40.
- Smith RR, Burke DA, Baldini AD, Shum-Siu A, Baltzley R, Bungler M, Magnuson DS. The Louisville Swim Scale: a novel assessment of hindlimb function following spinal cord injury in adult rats. *J Neurotrauma*. 2006 Nov;23(11):1654-70.
- Smith RR, Shum-Siu A, Baltzley R, Bungler M, Baldini A, Burke DA, Magnuson DS. Effects of swimming on functional recovery after incomplete spinal cord injury in rats. *J Neurotrauma*. 2006 Jun;23(6):908-19.
- Foldes ST, Taylor DM. Discreet discrete commands for assistive and neuroprosthetic devices. *IEEE Trans Neural Syst Rehabil Eng*. 2010 Jun;18(3):236-44.
- Marathe AR, Carey HL, Taylor DM. Virtual reality hardware and graphic display options for brain-machine interfaces. *J Neurosci Methods*. 2008 Jan 15;167(1):2-14. Epub 2007 Sep 29.
- Bardoni R, Takazawa T, Tong CK, Choudhury P, Scherrer G, Macdermott AB. Pre- and postsynaptic inhibitory control in the spinal cord dorsal horn. *Ann N Y Acad Sci*. 2013 Mar;1279:90-6.
- Shiokawa H, Kaftan EJ, MacDermott AB, Tong CK. NR2 subunits and NMDA receptors on lamina II inhibitory and excitatory interneurons of the mouse dorsal horn. *Mol Pain*. 2010 May 6;6:26.
- Tong CK, Kaftan EJ, Macdermott AB. Functional identification of NR2 subunits contributing to NMDA receptors on substance P receptor-expressing dorsal horn neurons. *Mol Pain*. 2008 Oct 10;4:44
- Tong CK, MacDermott AB. Both Ca²⁺-permeable and -impermeable AMPA receptors contribute to primary synaptic drive onto rat dorsal horn neurons. *J Physiol*. 2006 Aug 15;575(Pt 1):133-44
- Bardoni R, Torsney C, Tong CK, Prandini M, MacDermott AB. Presynaptic NMDA receptors modulate glutamate release from primary sensory neurons in rat spinal cord dorsal horn. *J Neurosci*. 2004 Mar 17;24(11):2774-81.
- Wang X, Arcuino G, Takano T, Lin J, Peng WG, Wan P, Li P, Xu Q, Liu QS, Goldman SA, Nedergaard M. P2X7 receptor inhibition improves recovery after spinal cord injury. *Nat Med*. 2004 Aug;10(8):821-7.

- Wang X, Lou N, Xu Q, Tian GF, Peng WG, Han X, Kang J, Takano T, Nedergaard M. Astrocytic Ca²⁺ signaling evoked by sensory stimulation in vivo. *Nat Neurosci*. 2006 Jun;9(6):816-23.
- Hook MA, Washburn SN, Moreno G, Woller SA, Puga D, Lee KH, Grau JW. An IL-1 receptor antagonist blocks a morphine-induced attenuation of locomotor recovery after spinal cord injury. *Brain Behav Immun*. 2011 Feb;25(2):349-59.
- Washburn SN, Maultsby ML, Puga DA, Grau JW. Opioid regulation of spinal cord plasticity: evidence the kappa-2 opioid receptor agonist GR89696 inhibits learning within the rat spinal cord. *Neurobiol Learn Mem*. 2008 Jan;89(1):1-16.
- Hook MA, Liu GT, Washburn SN, Ferguson AR, Bopp AC, Huie JR, Grau JW. The impact of morphine after a spinal cord injury. *Behav Brain Res*. 2007 May 16;179(2):281-93
- Washburn SN, Patton BC, Ferguson AR, Hudson KL, Grau JW. Exposure to intermittent nociceptive stimulation under pentobarbital anesthesia disrupts spinal cord function in rats. *Psychopharmacology (Berl)*. 2007 Jun;192(2):243-52
- Grau JW, Crown ED, Ferguson AR, Washburn SN, Hook MA, Miranda RC. Instrumental learning within the spinal cord: underlying mechanisms and implications for recovery after injury. *Behav Cogn Neurosci Rev*. 2006 Dec;5(4):191-239.
- Grau JW, Washburn SN, Hook MA, Ferguson AR, Crown ED, Garcia G, Bolding KA, Miranda RC. Uncontrollable stimulation undermines recovery after spinal cord injury. *J Neurotrauma*. 2004 Dec;21(12):1795-817.
- Ertürk A, Mauch CP, Hellal F, Förstner F, Keck T, Becker K, Jährling N, Steffens H, Richter M, Hübener M, Kramer E, Kirchhoff F, Dodt HU, Bradke F. Three-dimensional imaging of the unsectioned adult spinal cord to assess axon regeneration and glial responses after injury. *Nat Med*. 2011 Dec 25;18(1):166-71.
- Usher LC, Johnstone A, Ertürk A, Hu Y, Strikis D, Wanner IB, Moorman S, Lee JW, Min J, Ha HH, Duan Y, Hoffman S, Goldberg JL, Bradke F, Chang YT, Lemmon VP, Bixby JL. A chemical screen identifies novel compounds that overcome glial-mediated inhibition of neuronal regeneration. *J Neurosci*. 2010 Mar 31;30(13):4693-706.
- Cittelly DM, Nestic O, Johnson K, Hulsebosch C, Perez-Polo JR. Detrimental effects of antiapoptotic treatments in spinal cord injury. *Exp Neurol*. 2008 Apr;210(2):295-307
- Cittelly DM, Nestic-Taylor O, Perez-Polo JR. Phosphorylation of Bcl-xL after spinal cord injury. *J Neurosci Res*. 2007 Jul;85(9):1894-911.
- Nestic-Taylor O, Cittelly D, Ye Z, Xu GY, Unabia G, Lee JC, Svrakic NM, Liu XH, Youle RJ, Wood TG, McAdoo D, Westlund KN, Hulsebosch CE, Perez-Polo JR. Exogenous Bcl-xL fusion protein spares neurons after spinal cord injury. *J Neurosci Res*. 2005 Mar 1;79(5):628-37.

- Starkey ML, Bleul C, Maier IC, Schwab ME. Rehabilitative training following unilateral pyramidotomy in adult rats improves forelimb function in a non-task-specific way. *Exp Neurol*. 2011 Nov;232(1):81-9
- Maier IC, Ichiyama RM, Courtine G, Schnell L, Lavrov I, Edgerton VR, Schwab ME. Differential effects of anti-Nogo-A antibody treatment and treadmill training in rats with incomplete spinal cord injury. *Brain*. 2009 Jun;132(Pt 6):1426-40.
- Maier IC, Baumann K, Thallmair M, Weinmann O, Scholl J, Schwab ME. Constraint-induced movement therapy in the adult rat after unilateral corticospinal tract injury. *J Neurosci*. 2008 Sep 17;28(38):9386-403.
- Courtine G, Bunge MB, Fawcett JW, Grossman RG, Kaas JH, Lemon R, Maier I, Martin J, Nudo RJ, Ramon-Cueto A, Rouiller EM, Schnell L, Wannier T, Schwab ME, Edgerton VR. Can experiments in nonhuman primates expedite the translation of treatments for spinal cord injury in humans? *Nat Med*. 2007 May;13(5):561-6.
- Maier IC, Schwab ME. Sprouting, regeneration and circuit formation in the injured spinal cord: factors and activity. *Philos Trans R Soc Lond B Biol Sci*. 2006 Sep 29;361(1473):1611-34.
- Marino P, Norreel JC, Schachner M, Rougon G, Amoureux MC. A polysialic acid mimetic peptide promotes functional recovery in a mouse model of spinal cord injury. *Exp Neurol*. 2009 Sep;219(1):163-74.
- Nestic O, Sundberg LM, Herrera JJ, Mokkaapati VU, Lee J, Narayana PA. Vascular endothelial growth factor and spinal cord injury pain. *J Neurotrauma*. 2010 Oct;27(10):1793-803.
- Nestic O, Guest JD, Zivadinovic D, Narayana PA, Herrera JJ, Grill RJ, Mokkaapati VU, Gelman BB, Lee J. Aquaporins in spinal cord injury: the janus face of aquaporin 4. *Neuroscience*. 2010 Jul 28;168(4):1019-35.
- Nestic O, Lee J, Unabia GC, Johnson K, Ye Z, Vergara L, Hulsebosch CE, Perez-Polo JR. Aquaporin 1 - a novel player in spinal cord injury. *J Neurochem*. 2008 May;105(3):628-40.
- Nestic O, Lee J, Ye Z, Unabia GC, Rafati D, Hulsebosch CE, Perez-Polo JR. Acute and chronic changes in aquaporin 4 expression after spinal cord injury. *Neuroscience*. 2006 Dec;143(3):779-92.
- Nestic O, Lee J, Johnson KM, Ye Z, Xu GY, Unabia GC, Wood TG, McAdoo DJ, Westlund KN, Hulsebosch CE, Regino Perez-Polo J. Transcriptional profiling of spinal cord injury-induced central neuropathic pain. *J Neurochem*. 2005 Nov;95(4):998-1014.
- Sun YY, Wang CY, Hsu MF, Juan SH, Chang CY, Chou CM, Yang LY, Hung KS, Xu J, Lee YH, Hsu CY. Glucocorticoid protection of oligodendrocytes against excitotoxin involving hypoxia-inducible factor-1alpha in a cell-type-specific manner. *J Neurosci*. 2010 Jul 14;30(28):9621-30

- Lo WC, Hsu CH, Wu AT, Yang LY, Chen WH, Chiu WT, Lai WF, Wu CH, Gelovani JG, Deng WP. A novel cell-based therapy for contusion spinal cord injury using GDNF-delivering NIH3T3 cells with dual reporter genes monitored by molecular imaging. *J Nucl Med.* 2008 Sep;49(9):1512-9.
- Liu WL, Lee YH, Tsai SY, Hsu CY, Sun YY, Yang LY, Tsai SH, Yang WC. Methylprednisolone inhibits the expression of glial fibrillary acidic protein and chondroitin sulfate proteoglycans in reactivated astrocytes. *Glia.* 2008 Oct;56(13):1390-400.
- Tsai SY, Yang LY, Wu CH, Chang SF, Hsu CY, Wei CP, Leu SJ, Liaw J, Lee YH, Tsai MD. Injury-induced Janus kinase/protein kinase C-dependent phosphorylation of growth-associated protein 43 and signal transducer and activator of transcription 3 for neurite growth in dorsal root ganglion. *J Neurosci Res.* 2007 Feb 1;85(2):321-31
- Knerlich-Lukoschus F, Juraschek M, Blömer U, Lucius R, Mehdorn HM, Held-Feindt J. *J Neurotrauma.* 2008 May;25(5):427-48. Force-dependent development of neuropathic central pain and time-related CCL2/CCR2 expression after graded spinal cord contusion injuries of the rat.
- Afjehi-Sadat L, Brejnikow M, Kang SU, Vishwanath V, Walder N, Herkner K, Redl H, Lubec G. Differential protein levels and post-translational modifications in spinal cord injury of the rat. *J Proteome Res.* 2010 Mar 5;9(3):1591-7.
- Walder N, Petter-Puchner AH, Brejnikow M, Redl H, Essig M, Stieltjes B. Manganese enhanced magnetic resonance imaging in a contusion model of spinal cord injury in rats: correlation with motor function. *Invest Radiol.* 2008 May;43(5):277-83.
- Wang Y, Chen Y, Chen L, Wolpaw JR, Chen XY. Cortical stimulation causes long-term changes in H-reflexes and spinal motoneuron GABA receptors. *J Neurophysiol.* 2012 Nov;108(10):2668-78.
- Chen Y, Wang Y, Chen L, Sun C, English AW, Wolpaw JR, Chen XY. H-reflex up-conditioning encourages recovery of EMG activity and H-reflexes after sciatic nerve transection and repair in rats. *J Neurosci.* 2010 Dec 1;30(48):16128-36.
- Chen XY, Chen Y, Wang Y, Thompson A, Carp JS, Segal RL, Wolpaw JR. bReflex conditioning: a new strategy for improving motor function after spinal cord injury. *Ann N Y Acad Sci.* 2010 Jun;1198 Suppl 1:E12-21.
- Chen XY, Pillai S, Chen Y, Wang Y, Chen L, Carp JS, Wolpaw JR. Spinal and supraspinal effects of long-term stimulation of sensorimotor cortex in rats. *J Neurophysiol.* 2007 Aug;98(2):878-87.
- Chen Y, Chen XY, Jakeman LB, Chen L, Stokes BT, Wolpaw JR. Operant conditioning of H-reflex can correct a locomotor abnormality after spinal cord injury in rats. *J Neurosci.* 2006 Nov 29;26(48):12537-43.

- Chen XY, Chen Y, Chen L, Tennissen AM, Wolpaw JR Corticospinal tract transection permanently abolishes H-reflex down-conditioning in rats. *J Neurotrauma*. 2006 Nov;23(11):1705-12.
- Chen XY, Chen L, Chen Y, Wolpaw JR. Operant conditioning of reciprocal inhibition in rat soleus muscle. *J Neurophysiol*. 2006 Oct;96(4):2144-50.
- Chen Y, Chen XY, Jakeman LB, Schalk G, Stokes BT, Wolpaw JR. The interaction of a new motor skill and an old one: H-reflex conditioning and locomotion in rats. *J Neurosci*. 2005 Jul 20;25(29):6898-906.
- Fouts DE, Pieper R, Szpakowski S, Pohl H, Knoblach S, Suh MJ, Huang ST, Ljungberg I, Sprague BM, Lucas SK, Torralba M, Nelson KE, Groah SL. Integrated next-generation sequencing of 16S rDNA and metaproteomics differentiate the healthy urine microbiome from asymptomatic bacteriuria in neuropathic bladder associated with spinal cord injury. *J Transl Med*. 2012 Aug 28;10:174.
- Pajoohesh-Ganji A, Knoblach SM, Faden AI, Byrnes KR. Characterization of inflammatory gene expression and galectin-3 function after spinal cord injury in mice. *Brain Res*. 2012 Sep 26;1475:96-105.
- Byrnes KR, Washington PM, Knoblach SM, Hoffman E, Faden AI. Delayed inflammatory mRNA and protein expression after spinal cord injury. *J Neuroinflammation*. 2011 Oct 5;8:130.
- Kim B, Lee B, Knoblach S, Hoffman E, Seo J. GeneShelf: a web-based visual interface for large gene expression time-series data repositories. *IEEE Trans Vis Comput Graph*. 2009 Nov-Dec;15(6):905-12.
- Byrnes KR, Garay J, Di Giovanni S, De Biase A, Knoblach SM, Hoffman EP, Movsesyan V, Faden AI. Expression of two temporally distinct microglia-related gene clusters after spinal cord injury. *Glia*. 2006 Mar;53(4):420-33.
- De Biase A, Knoblach SM, Di Giovanni S, Fan C, Molon A, Hoffman EP, Faden AI. Gene expression profiling of experimental traumatic spinal cord injury as a function of distance from impact site and injury severity. *Physiol Genomics*. 2005 Aug 11;22(3):368-81.
- Faden AI, Movsesyan VA, Knoblach SM, Ahmed F, Cernak I. Neuroprotective effects of novel small peptides in vitro and after brain injury. *Neuropharmacology*. 2005 Sep;49(3):410-24.
- Knoblach SM, Huang X, VanGelderen J, Calva-Cerqueira D, Faden AI. Selective caspase activation may contribute to neurological dysfunction after experimental spinal cord trauma. *J Neurosci Res*. 2005 May 1;80(3):369-80.
- Caggiano AO, Zimmer MP, Ganguly A, Blight AR, Gruskin EA. Chondroitinase ABCI improves locomotion and bladder function following contusion injury of the rat spinal cord. *J Neurotrauma*. 2005 Feb;22(2):226-39.

- Li Y, Oskouian RJ, Day YJ, Rieger JM, Liu L, Kern JA, Linden J. Mouse spinal cord compression injury is reduced by either activation of the adenosine A2A receptor on bone marrow-derived cells or deletion of the A2A receptor on non-bone marrow-derived cells. *Neuroscience*. 2006 Sep 15;141(4):2029-39
- Li Y, Oskouian RJ, Day YJ, Kern JA, Linden J. Optimization of a mouse locomotor rating system to evaluate compression-induced spinal cord injury: correlation of locomotor and morphological injury indices. *J Neurosurg Spine*. 2006 Feb;4(2):165-73.
- Agudo M, Yip P, Davies M, Bradbury E, Doherty P, McMahon S, Maden M, Corcoran JP. A retinoic acid receptor beta agonist (CD2019) overcomes inhibition of axonal outgrowth via phosphoinositide 3-kinase signalling in the injured adult spinal cord. *Neurobiol Dis*. 2010 Jan;37(1):147-55.
- Starkey ML, Davies M, Yip PK, Carter LM, Wong DJ, McMahon SB, Bradbury EJ. Expression of the regeneration-associated protein SPRR1A in primary sensory neurons and spinal cord of the adult mouse following peripheral and central injury. *J Comp Neurol*. 2009 Mar 1;513(1):51-68.
- Carter LM, Starkey ML, Akrimi SF, Davies M, McMahon SB, Bradbury EJ. The yellow fluorescent protein (YFP-H) mouse reveals neuroprotection as a novel mechanism underlying chondroitinase ABC-mediated repair after spinal cord injury. *J Neurosci*. 2008 Dec 24;28(52):14107-20.
- Barritt AW, Davies M, Marchand F, Hartley R, Grist J, Yip P, McMahon SB, Bradbury EJ. Chondroitinase ABC promotes sprouting of intact and injured spinal systems after spinal cord injury. *J Neurosci*. 2006 Oct 18;26(42):10856-67.
- Starkey ML, Barritt AW, Yip PK, Davies M, Hamers FP, McMahon SB, Bradbury EJ. Assessing behavioural function following a pyramidotomy lesion of the corticospinal tract in adult mice. *Exp Neurol*. 2005 Oct;195(2):524-39.
- Curinga G, Smith GM. Molecular/genetic manipulation of extrinsic axon guidance factors for CNS repair and regeneration. *Exp Neurol*. 2008 Feb;209(2):333-42.
- Curinga GM, Snow DM, Mashburn C, Kohler K, Thobaben R, Caggiano AO, Smith GM. Mammalian-produced chondroitinase AC mitigates axon inhibition by chondroitin sulfate proteoglycans. *J Neurochem*. 2007 Jul;102(1):275-8805
- Fortun J, Hill CE, Bunge MB. Combinatorial strategies with Schwann cell transplantation to improve repair of the injured spinal cord. *Neurosci Lett*. 2009 Jun 12;456(3):124-32.
- Fortun J, Puzis R, Pearse DD, Gage FH, Bunge MB. Muscle injection of AAV-NT3 promotes anatomical reorganization of CST axons and improves behavioral outcome following SCI. *J Neurotrauma*. 2009 Jul;26(7):941-53

- Gonzenbach RR, Zoerner B, Schnell L, Weinmann O, Mir AK, Schwab ME. Delayed anti-nogo-a antibody application after spinal cord injury shows progressive loss of responsiveness. *J Neurotrauma*. 2012 Feb 10;29(3):567-78.
- Gonzenbach RR, Gasser P, Zörner B, Hochreutener E, Dietz V, Schwab ME. Nogo-A antibodies and training reduce muscle spasms in spinal cord-injured rats. *Ann Neurol*. 2010 Jul;68(1):48-57.
- Müllner A, Gonzenbach RR, Weinmann O, Schnell L, Liebscher T, Schwab ME. Lamina-specific restoration of serotonergic projections after Nogo-A antibody treatment of spinal cord injury in rats. *Eur J Neurosci*. 2008 Jan;27(2):326-33.
- Gonzenbach RR, Schwab ME. Disinhibition of neurite growth to repair the injured adult CNS: focusing on Nogo. *Cell Mol Life Sci*. 2008 Jan;65(1):161-76.
- Brewer KL, Nolan TA. Spinal and supraspinal changes in tumor necrosis factor-alpha expression following excitotoxic spinal cord injury. *J Mol Neurosci*. 2007;31(1):13-21.
- Britt JM, Kane JR, Spaeth CS, Zuzek A, Robinson GL, Gbanaglo MY, Estler CJ, Boydston EA, Schallert T, Bittner GD. Polyethylene glycol rapidly restores axonal integrity and improves the rate of motor behavior recovery after sciatic nerve crush injury. *J Neurophysiol*. 2010 Aug;104(2):695-703.
- Duffy P, Wang X, Siegel CS, Tu N, Henkemeyer M, Cafferty WB, Strittmatter SM. Myelin-derived ephrinB3 restricts axonal regeneration and recovery after adult CNS injury. *Proc Natl Acad Sci U S A*. 2012 Mar 27;109(13):5063-8.
- Wang X, Duffy P, McGee AW, Hasan O, Gould G, Tu N, Harel NY, Huang Y, Carson RE, Weinzimmer D, Ropchan J, Benowitz LI, Cafferty WB, Strittmatter SM. Recovery from chronic spinal cord contusion after Nogo receptor intervention. *Ann Neurol*. 2011 Nov;70(5):805-21.
- Cafferty WB, Duffy P, Huebner E, Strittmatter SM. MAG and OMgp synergize with Nogo-A to restrict axonal growth and neurological recovery after spinal cord trauma. *J Neurosci*. 2010 May 19;30(20):6825-37.
- Duffy P, Schmandke A, Schmandke A, Sigworth J, Narumiya S, Cafferty WB, Strittmatter SM. Rho-associated kinase II (ROCKII) limits axonal growth after trauma within the adult mouse spinal cord. *J Neurosci*. 2009 Dec 2;29(48):15266-76
- Cafferty WB, Bradbury EJ, Lidieth M, Jones M, Duffy PJ, Pezet S, McMahon SB. Chondroitinase ABC-mediated plasticity of spinal sensory function. *J Neurosci*. 2008 Nov 12;28(46):11998-2009.
- Cafferty WB, Yang SH, Duffy PJ, Li S, Strittmatter SM. Functional axonal regeneration through astrocytic scar genetically modified to digest chondroitin sulfate proteoglycans. *J Neurosci*. 2007 Feb 28;27(9):2176-85.

- Liu J, Akay T, Hedlund PB, Pearson KG, Jordan LM. Spinal 5-HT7 receptors are critical for alternating activity during locomotion: in vitro neonatal and in vivo adult studies using 5-HT7 receptor knockout mice. *J Neurophysiol.* 2009 Jul;102(1):337-48.
- Carlin KP, Liu J, Jordan LM. Postnatal changes in the inactivation properties of voltage-gated sodium channels contribute to the mature firing pattern of spinal motoneurons. *J Neurophysiol.* 2008 Jun;99(6):2864-76.
- Jordan LM, Liu J, Hedlund PB, Akay T, Pearson KG. Descending command systems for the initiation of locomotion in mammals. *Brain Res Rev.* 2008 Jan;57(1):183-91.
- Liu J, Jordan LM. Stimulation of the parapyramidal region of the neonatal rat brain stem produces locomotor-like activity involving spinal 5-HT7 and 5-HT2A receptors. *J Neurophysiol.* 2005 Aug;94(2):1392-404.
- Kim JH, Song SK. Diffusion tensor imaging of the mouse brainstem and cervical spinal cord. *Nat Protoc.* 2013 Feb;8(2):409-17.
- Kim JH, Song SK, Burke DA, Magnuson DS. Comprehensive locomotor outcomes correlate to hyperacute diffusion tensor measures after spinal cord injury in the adult rat. *Exp Neurol.* 2012 May;235(1):188-96.
- Reddy AS, Kim JH, Hawkins-Salsbury JA, Macauley SL, Tracy ET, Vogler CA, Han X, Song SK, Wozniak DF, Fowler SC, Klein RS, Sands MS. Bone marrow transplantation augments the effect of brain- and spinal cord-directed adeno-associated virus 2/5 gene therapy by altering inflammation in the murine model of globoid-cell leukodystrophy. *J Neurosci.* 2011 Jul 6;31(27):9945-57.
- Kim JH, Wu TH, Budde MD, Lee JM, Song SK. Noninvasive detection of brainstem and spinal cord axonal degeneration in an amyotrophic lateral sclerosis mouse model. *NMR Biomed.* 2011 Feb;24(2):163-9.
- Tidwell VK, Kim JH, Song SK, Nehorai A. Automatic segmentation of rodent spinal cord diffusion MR images. *Magn Reson Med.* 2010 Sep;64(3):893-901.
- Kim JH, Loy DN, Wang Q, Budde MD, Schmidt RE, Trinkaus K, Song SK. Diffusion tensor imaging at 3 hours after traumatic spinal cord injury predicts long-term locomotor recovery. *J Neurotrauma.* 2010 Mar;27(3):587-98.
- Tu TW, Kim JH, Wang J, Song SK. Full tensor diffusion imaging is not required to assess the white-matter integrity in mouse contusion spinal cord injury. *J Neurotrauma.* 2010 Jan;27(1):253-62.
- Hofling AA, Kim JH, Fantz CR, Sands MS, Song SK. Diffusion tensor imaging detects axonal injury and demyelination in the spinal cord and cranial nerves of a murine model of globoid cell leukodystrophy. *NMR Biomed.* 2009 Dec;22(10):1100-6.

- Kim JH, Tu TW, Bayly PV, Song SK. Impact speed does not determine severity of spinal cord injury in mice with fixed impact displacement. *J Neurotrauma*. 2009 Aug;26(8):1395-404
- Kim JH, Haldar J, Liang ZP, Song SK. Diffusion tensor imaging of mouse brain stem and cervical spinal cord. *J Neurosci Methods*. 2009 Jan 30;176(2):186-91.
- Budde MD, Kim JH, Liang HF, Russell JH, Cross AH, Song SK. Axonal injury detected by in vivo diffusion tensor imaging correlates with neurological disability in a mouse model of multiple sclerosis. *NMR Biomed*. 2008 Jul;21(6):589-97.
- Kim JH, Loy DN, Liang HF, Trinkaus K, Schmidt RE, Song SK. Noninvasive diffusion tensor imaging of evolving white matter pathology in a mouse model of acute spinal cord injury. *Magn Reson Med*. 2007 Aug;58(2):253-60.
- Loy DN, Kim JH, Xie M, Schmidt RE, Trinkaus K, Song SK. Diffusion tensor imaging predicts hyperacute spinal cord injury severity. *J Neurotrauma*. 2007 Jun;24(6):979-90.
- Kim JH, Trinkaus K, Ozcan A, Budde MD, Song SK. Postmortem delay does not change regional diffusion anisotropy characteristics in mouse spinal cord white matter. *NMR Biomed*. 2007 May;20(3):352-9.
- Budde MD, Kim JH, Liang HF, Schmidt RE, Russell JH, Cross AH, Song SK. Toward accurate diagnosis of white matter pathology using diffusion tensor imaging. *Magn Reson Med*. 2007 Apr;57(4):688-95.
- Kim JH, Budde MD, Liang HF, Klein RS, Russell JH, Cross AH, Song SK. Detecting axon damage in spinal cord from a mouse model of multiple sclerosis. *Neurobiol Dis*. 2006 Mar;21(3):626-32.
- Schroeder JL, Highsmith JM, Young HF, Mathern BE. Reduction of hypoxia by perfluorocarbon emulsion in a traumatic spinal cord injury model. *J Neurosurg Spine*. 2008 Aug;9(2):213-20.
- Marques SA, Garcez VF, Del Bel EA, Martinez AM. A simple, inexpensive and easily reproducible model of spinal cord injury in mice: morphological and functional assessment. *J Neurosci Methods*. 2009 Feb 15;177(1):183-93.
- Mountney A, Zahner MR, Sturgill ER, Riley CJ, Aston JW, Oudega M, Schramm LP, Hurtado A, Schnaar RL. Sialidase, chondroitinase ABC, and combination therapy after spinal cord contusion injury. *J Neurotrauma*. 2013 Feb 1;30(3):181-90.
- Benedict AL, Mountney A, Hurtado A, Bryan KE, Schnaar RL, Dinkova-Kostova AT, Talalay P. Neuroprotective effects of sulforaphane after contusive spinal cord injury. *J Neurotrauma*. 2012 Nov 1;29(16):2576-86

- Mountney A, Zahner MR, Lorenzini I, Oudega M, Schramm LP, Schnaar RL. Sialidase enhances recovery from spinal cord contusion injury. *Proc Natl Acad Sci U S A*. 2010 Jun 22;107(25):11561-6.
- Yang LJ, Lorenzini I, Vajn K, Mountney A, Schramm LP, Schnaar RL. Sialidase enhances spinal axon outgrowth in vivo. *Proc Natl Acad Sci U S A*. 2006 Jul 18;103(29):11057-62.
- Bottai D, Cigognini D, Madaschi L, Adami R, Nicora E, Menarini M, Di Giulio AM, Gorio A. Embryonic stem cells promote motor recovery and affect inflammatory cell infiltration in spinal cord injured mice. *Exp Neurol*. 2010 Jun;223(2):452-63.
- Bottai D, Madaschi L, Di Giulio AM, Gorio A. Viability-dependent promoting action of adult neural precursors in spinal cord injury. *Mol Med*. 2008 Sep-Oct;14(9-10):634-44
- Zhang X, Douglas KL, Jin H, Eldaif BM, Nassar R, Fraser MO, Dolber PC. Sprouting of substance P-expressing primary afferent central terminals and spinal micturition reflex NK1 receptor dependence after spinal cord injury. *Am J Physiol Regul Integr Comp Physiol*. 2008 Dec;295(6):R2084-96.
- Gu B, Thor KB, Reiter JP, Dolber PC. Effect of 5-hydroxytryptamine₁ serotonin receptor agonists on noxiously stimulated micturition in cats with chronic spinal cord injury. *J Urol*. 2007 Jun;177(6):2381-5.
- Pandit V, Zuidema J, Venuto KN, Macione J, Dai G, Gilbert RJ, Kotha S. Evaluation of Multifunctional Polysaccharide Hydrogels with Varying Stiffness for Bone Tissue Engineering. *Tissue Eng Part A*. 2013 Jun 2.
- Hurtado A, Cregg JM, Wang HB, Wendell DF, Oudega M, Gilbert RJ, McDonald JW. Robust CNS regeneration after complete spinal cord transection using aligned poly-L-lactic acid microfibers. *Biomaterials*. 2011 Sep;32(26):6068-79
- Gilbert RJ, Rivet CJ, Zuidema JM, Popovich PG. Biomaterial design considerations for repairing the injured spinal cord. *Crit Rev Biomed Eng*. 2011;39(2):125-80.
- Hellal F, Hurtado A, Ruschel J, Flynn KC, Laskowski CJ, Umlauf M, Kapitein LC, Strikis D, Lemmon V, Bixby J, Hoogenraad CC, Bradke F. Microtubule stabilization reduces scarring and causes axon regeneration after spinal cord injury. *Science*. 2011 Feb 18;331(6019):928-31.
- Arevalo-Martin A, Garcia-Ovejero D, Sierra-Palomares Y, Paniagua-Torija B, Gonzalez-Gil I, Ortega-Gutierrez S, Molina-Holgado E. Early endogenous activation of CB1 and CB2 receptors after spinal cord injury is a protective response involved in spontaneous recovery. *PLoS One*. 2012;7(11):e49057.

- Garcia-Ovejero D, Arevalo-Martin A, Paniagua-Torija B, Sierra-Palomares Y, Molina-Holgado E. A cell population that strongly expresses the CB1 cannabinoid receptor in the ependyma of the rat spinal cord. *J Comp Neurol*. 2013 Jan 1;521(1):233-51.
- Oliviero A, Arevalo-Martin A, Rotondi M, García-Ovejero D, Mordillo-Mateos L, Lozano-Sicilia A, Panyavin I, Chiovato L, Aguilar J, Foffani G, Di Lazzaro V, Molina-Holgado E. CB1 receptor antagonism/inverse agonism increases motor system excitability in humans. *Eur Neuropsychopharmacol*. 2012 Jan;22(1):27-35.
- Arevalo-Martin A, Garcia-Ovejero D, Molina-Holgado E. The endocannabinoid 2-arachidonoylglycerol reduces lesion expansion and white matter damage after spinal cord injury. *Neurobiol Dis*. 2010 May;38(2):304-12.
- Garcia-Ovejero D, Arevalo-Martin A, Petrosino S, Docagne F, Hagen C, Bisogno T, Watanabe M, Guaza C, Di Marzo V, Molina-Holgado E. The endocannabinoid system is modulated in response to spinal cord injury in rats. *Neurobiol Dis*. 2009 Jan;33(1):57-71.
- Pal R, Gopinath C, Rao NM, Banerjee P, Krishnamoorthy V, Venkataramana NK, Totey S. Functional recovery after transplantation of bone marrow-derived human mesenchymal stromal cells in a rat model of spinal cord injury. *Cytotherapy*. 2010 Oct;12(6):792-806.
- Pal R, Venkataramana NK, Bansal A, Balaraju S, Jan M, Chandra R, Dixit A, Rauthan A, Murgod U, Totey S. Ex vivo-expanded autologous bone marrow-derived mesenchymal stromal cells in human spinal cord injury/paraplegia: a pilot clinical study. *Cytotherapy*. 2009;11(7):897-911.
- White TE, Lane MA, Sandhu MS, O'Steen BE, Fuller DD, Reier PJ. Neuronal progenitor transplantation and respiratory outcomes following upper cervical spinal cord injury in adult rats. *Exp Neurol*. 2010 Sep;225(1):231-6.
- Lee KZ, Sandhu MS, Dougherty BJ, Reier PJ, Fuller DD. Influence of vagal afferents on supraspinal and spinal respiratory activity following cervical spinal cord injury in rats. *J Appl Physiol*. 2010 Aug;109(2):377-87.
- Sandhu MS, Dougherty BJ, Lane MA, Bolser DC, Kirkwood PA, Reier PJ, Fuller DD. Respiratory recovery following high cervical hemisection. *Respir Physiol Neurobiol*. 2009 Nov 30;169(2):94-101.
- Fuller DD, Sandhu MS, Doperalski NJ, Lane MA, White TE, Bishop MD, Reier PJ. Graded unilateral cervical spinal cord injury and respiratory motor recovery. *Respir Physiol Neurobiol*. 2009 Feb 28;165(2-3):245-53.
- Lane MA, White TE, Coutts MA, Jones AL, Sandhu MS, Bloom DC, Bolser DC, Yates BJ, Fuller DD, Reier PJ. Cervical prephrenic interneurons in the normal and lesioned spinal cord of the adult rat. *J Comp Neurol*. 2008 Dec 10;511(5):692-709.

- Doperalski NJ, Sandhu MS, Bavis RW, Reier PJ, Fuller DD. Ventilation and phrenic output following high cervical spinal hemisection in male vs. female rats. *Respir Physiol Neurobiol.* 2008 Jul 31;162(2):160-7.
- Fuller DD, Doperalski NJ, Dougherty BJ, Sandhu MS, Bolser DC, Reier PJ. Modest spontaneous recovery of ventilation following chronic high cervical hemisection in rats. *Exp Neurol.* 2008 May;211(1):97-106.
- Harvey P, Gong B, Rossomando AJ, Frank E. Topographically specific regeneration of sensory axons in the spinal cord. *Proc Natl Acad Sci U S A.* 2010 Jun 22;107(25):11585-90.
- Harvey PA, Lee DH, Qian F, Weinreb PH, Frank E. Blockade of Nogo receptor ligands promotes functional regeneration of sensory axons after dorsal root crush. *J Neurosci.* 2009 May 13;29(19):6285-95.
- Wang R, King T, Ossipov MH, Rossomando AJ, Vanderah TW, Harvey P, Cariani P, Frank E, Sah DW, Porreca F. Persistent restoration of sensory function by immediate or delayed systemic artemin after dorsal root injury. *Nat Neurosci.* 2008 Apr;11(4):488-96.
- Zörner B, Filli L, Starkey ML, Gonzenbach R, Kasper H, Röthlisberger M, Bolliger M, Schwab ME. Profiling locomotor recovery: comprehensive quantification of impairments after CNS damage in rodents. *Nat Methods.* 2010 Sep;7(9):701-8.
- Zörner B, Schwab ME. Anti-Nogo on the go: from animal models to a clinical trial. *Ann N Y Acad Sci.* 2010 Jun;1198 Suppl 1:E22-34.
- Obermair FJ, Fiorelli R, Schroeter A, Beyeler S, Blatti C, Zoerner B, Thallmair M. A novel classification of quiescent and transit amplifying adult neural stem cells by surface and metabolic markers permits a defined simultaneous isolation. *Stem Cell Res.* 2010 Sep;5(2):131-43.
- Zörner B, Blanckenhorn WU, Dietz V; EM-SCI Study Group, Curt A. Clinical algorithm for improved prediction of ambulation and patient stratification after incomplete spinal cord injury. *J Neurotrauma.* 2010 Jan;27(1):241-52.
- Yunta M, Nieto-Díaz M, Esteban FJ, Caballero-López M, Navarro-Ruíz R, Reigada D, Pita-Thomas DW, del Águila A, Muñoz-Galdeano T, Maza RM. MicroRNA dysregulation in the spinal cord following traumatic injury. *PLoS One.* 2012;7(4):e34534.
- Carelli S, Marfia G, Di Giulio AM, Ghilardi G, Gorio A. Erythropoietin: recent developments in the treatment of spinal cord injury. *Neurol Res Int.* 2011;2011:453179.
- Fumagalli F, Madaschi L, Caffino L, Marfia G, Di Giulio AM, Racagni G, Gorio A. Acute spinal cord injury reduces brain derived neurotrophic factor expression in rat hippocampus. *Neuroscience.* 2009 Mar 31;159(3):936-9

- Fumagalli F, Madaschi L, Brenna P, Caffino L, Marfia G, Di Giulio AM, Racagni G, Gorio A. Single exposure to erythropoietin modulates Nerve Growth Factor expression in the spinal cord following traumatic injury: comparison with methylprednisolone. *Eur J Pharmacol.* 2008 Jan 6;578(1):19-27.
- Saxena T, Gilbert J, Stelzner D, Hasenwinkel J. Mechanical characterization of the injured spinal cord after lateral spinal hemisection injury in the rat. *J Neurotrauma.* 2012 Jun 10;29(9):1747-57.
- Saxena T, Deng B, Stelzner D, Hasenwinkel J, Chaiken J. Raman spectroscopic investigation of spinal cord injury in a rat model. *J Biomed Opt.* 2011 Feb;16(2):027003.
- Floriddia EM, Rathore KI, Tedeschi A, Quadrato G, Wuttke A, Lueckmann JM, Kigerl KA, Popovich PG, Di Giovanni S. p53 Regulates the neuronal intrinsic and extrinsic responses affecting the recovery of motor function following spinal cord injury. *J Neurosci.* 2012 Oct 3;32(40):13956-70
- Liu K, Lu Y, Lee JK, Samara R, Willenberg R, Sears-Kraxberger I, Tedeschi A, Park KK, Jin D, Cai B, Xu B, Connolly L, Steward O, Zheng B, He Z. PTEN deletion enhances the regenerative ability of adult corticospinal neurons. *Nat Neurosci.* 2010 Sep;13(9):1075-81.
- Tedeschi A, Nguyen T, Steele SU, Feil S, Naumann U, Feil R, Di Giovanni S. The tumor suppressor p53 transcriptionally regulates cGKI expression during neuronal maturation and is required for cGMP-dependent growth cone collapse. *J Neurosci.* 2009 Dec 2;29(48):15155-60.
- Liu K, Lu Y, Lee JK, Samara R, Willenberg R, Sears-Kraxberger I, Tedeschi A, Park KK, Jin D, Cai B, Xu B, Connolly L, Steward O, Zheng B, He Z. PTEN deletion enhances the regenerative ability of adult corticospinal neurons. *Nat Neurosci.* 2010 Sep;13(9):1075-81
- Liu K, Tedeschi A, Park KK, He Z. Neuronal intrinsic mechanisms of axon regeneration. *Annu Rev Neurosci.* 2011;34:131-52.
- Tedeschi A, He Z. Axon regeneration: electrical silencing is a condition for regrowth. *Curr Biol.*
- Tedeschi A. Tuning the orchestra: transcriptional pathways controlling axon regeneration. *Front Mol Neurosci.* 2011;4:60.
- DeRuisseau LR, Recca DM, Mogle JA, Zoccolillo M, DeRuisseau KC. Metallothionein deficiency leads to soleus muscle contractile dysfunction following acute spinal cord injury in mice. *Am J Physiol Regul Integr Comp Physiol.* 2009 Dec;297(6):R1795-802.
- Huang C, Han X, Li X, Lam E, Peng W, Lou N, Torres A, Yang M, Garre JM, Tian GF, Bennett MV, Nedergaard M, Takano T. Critical role of connexin 43 in secondary expansion of traumatic spinal cord injury. *J Neurosci.* 2012 Mar 7;32(10):3333-8

- Peng W, Cotrina ML, Han X, Yu H, Bekar L, Blum L, Takano T, Tian GF, Goldman SA, Nedergaard M. Systemic administration of an antagonist of the ATP-sensitive receptor P2X7 improves recovery after spinal cord injury. *Proc Natl Acad Sci U S A*. 2009 Jul 28;106(30):12489-93.
- Guo X, Das M, Rumsey J, Gonzalez M, Stancescu M, Hickman J. Neuromuscular junction formation between human stem-cell-derived motoneurons and rat skeletal muscle in a defined system. *Tissue Eng Part C Methods*. 2010 Dec;16(6):1347-55.
- Baumbauer KM, Lee KH, Puga DA, Woller SA, Hughes AJ, Grau JW. Temporal regularity determines the impact of electrical stimulation on tactile reactivity and response to capsaicin in spinally transected rats. *Neuroscience*. 2012 Dec 27;227:119-33.
- Hook MA, Moreno G, Woller S, Puga D, Hoy K Jr, Balden R, Grau JW. Intrathecal morphine attenuates recovery of function after a spinal cord injury. *J Neurotrauma*. 2009 May;26(5):741-52
- Baumbauer KM, Hoy KC Jr, Huie JR, Hughes AJ, Woller SA, Puga DA, Setlow B, Grau JW. Timing in the absence of supraspinal input I: variable, but not fixed, spaced stimulation of the sciatic nerve undermines spinally-mediated instrumental learning. *Neuroscience*. 2008 Sep 9;155(4):1030-47.
- Wang H, Katagiri Y, McCann TE, Unsworth E, Goldsmith P, Yu ZX, Tan F, Santiago L, Mills EM, Wang Y, Symes AJ, Geller HM. Chondroitin-4-sulfation negatively regulates axonal guidance and growth. *J Cell Sci*. 2008 Sep 15;121(Pt 18):3083-91.
- Johnstone JT, Morton PD, Jayakumar AR, Bracchi-Ricard V, Runko E, Liebl DJ, Norenberg MD, Bethea JR. Reduced extracellular zinc levels facilitate glutamate-mediated oligodendrocyte death after trauma. *J Neurosci Res*. 2013 Jun;91(6):828-37.
- Morton PD, Johnstone JT, Ramos AY, Liebl DJ, Bunge MB, Bethea JR. Nuclear factor- κ B activation in Schwann cells regulates regeneration and remyelination. *Glia*. 2012 Apr;60(4):639-50.
- Zhang H, Trivedi A, Lee JU, Lohela M, Lee SM, Fandel TM, Werb Z, Noble-Haeusslein LJ. Matrix metalloproteinase-9 and stromal cell-derived factor-1 act synergistically to support migration of blood-borne monocytes into the injured spinal cord. *J Neurosci*. 2011 Nov 2;31(44):15894-903
- Lee SM, Rosen S, Weinstein P, van Rooijen N, Noble-Haeusslein LJ. Prevention of both neutrophil and monocyte recruitment promotes recovery after spinal cord injury. *J Neurotrauma*. 2011 Sep;28(9):1893-907.
- Almad A, Lash AT, Wei P, Lovett-Racke AE, McTigue DM. The PPAR alpha agonist gemfibrozil is an ineffective treatment for spinal cord injured mice. *Exp Neurol*. 2011 Dec;232(2):309-17
- Almad A, Sahinkaya FR, McTigue DM. Oligodendrocyte fate after spinal cord injury. *Neurotherapeutics*. 2011 Apr;8(2):262-73.

- Almad A, McTigue DM. Chronic expression of PPAR-delta by oligodendrocyte lineage cells in the injured rat spinal cord. *J Comp Neurol*. 2010 Mar 15;518(6):785-99.
- Gensel JC, Almad AA, Alexander JK, Schonberg DL, Tripathi RB. Does chronic remyelination occur for all spared axons after spinal cord injury in mouse? *J Neurosci*. 2008 Aug 20;28(34):8385-6.
- Porensky PN, Mitrapant C, McGovern VL, Bevan AK, Foust KD, Kaspar BK, Wilton SD, Burghes AH. A single administration of morpholino antisense oligomer rescues spinal muscular atrophy in mouse. *Hum Mol Genet*. 2012 Apr 1;21(7):1625-38.
- Bevan AK, Duque S, Foust KD, Morales PR, Braun L, Schmelzer L, Chan CM, McCrate M, Chicoine LG, Coley BD, Porensky PN, Kolb SJ, Mendell JR, Burghes AH, Kaspar BK. Systemic gene delivery in large species for targeting spinal cord, brain, and peripheral tissues for pediatric disorders. *Mol Ther*. 2011 Nov;19(11):1971-80.
- Dulin JN, Karoly ED, Wang Y, Strobel HW, Grill RJ. Licofelone modulates neuroinflammation and attenuates mechanical hypersensitivity in the chronic phase of spinal cord injury. *J Neurosci*. 2013 Jan 9;33(2):652-64.
- Dulin JN, Moore ML, Grill RJ. The dual cyclooxygenase/5-lipoxygenase inhibitor licofelone attenuates p-glycoprotein-mediated drug resistance in the injured spinal cord. *J Neurotrauma*. 2013 Feb 1;30(3):211-26.
- Dulin JN, Moore ML, Gates KW, Queen JH, Grill RJ. Spinal cord injury causes sustained disruption of the blood-testis barrier in the rat. *PLoS One*. 2011 Jan 26;6(1):e16456.
- Streijger F, Beernink TM, Lee JH, Bhatnagar T, Park S, Kwon BK, Tetzlaff W. Characterization of a cervical spinal cord hemicontusion injury in mice using the infinite horizon impactor. *J Neurotrauma*. 2013 May 15;30(10):869-83.
- Lee JH, Streijger F, Tigchelaar S, Maloon M, Liu J, Tetzlaff W, Kwon BK. A contusive model of unilateral cervical spinal cord injury using the infinite horizon impactor. *J Vis Exp*. 2012 Jul 24;(65)
- Lubieniecka JM, Streijger F, Lee JH, Stoyanov N, Liu J, Mottus R, Pfeifer T, Kwon BK, Coorsen JR, Foster LJ, Grigliatti TA, Tetzlaff W. Biomarkers for severity of spinal cord injury in the cerebrospinal fluid of rats. *PLoS One*. 2011 Apr 29;6(4):e19247.
- Streijger F, Plunet WT, Plemel JR, Lam CK, Liu J, Tetzlaff W. Intermittent fasting in mice does not improve hindlimb motor performance after spinal cord injury. *J Neurotrauma*. 2011 Jun;28(6):1051-61.
- Jeong MA, Plunet W, Streijger F, Lee JH, Plemel JR, Park S, Lam CK, Liu J, Tetzlaff W. Intermittent fasting improves functional recovery after rat thoracic contusion spinal cord injury. *J Neurotrauma*. 2011 Mar;28(3):479-92. doi: 10.1089/neu.2010.1609.

- Lee JH, Tigchelaar S, Liu J, Stammers AM, Streijger F, Tetzlaff W, Kwon BK. Lack of neuroprotective effects of simvastatin and minocycline in a model of cervical spinal cord injury. *Exp Neurol*. 2010 ep;225(1):219-30.
- Hawthorne AL, Hu H, Kundu B, Steinmetz MP, Wylie CJ, Deneris ES, Silver J. The unusual response of serotonergic neurons after CNS Injury: lack of axonal dieback and enhanced sprouting within the inhibitory environment of the glial scar. *J Neurosci*. 2011 Apr 13;31(15):5605-16.
- Hawthorne AL, Popovich PG. Emerging concepts in myeloid cell biology after spinal cord injury. *Neurotherapeutics*. 2011 Apr;8(2):252-61.
- Busch SA, Horn KP, Cuascut FX, Hawthorne AL, Bai L, Miller RH, Silver J. Adult NG2+ cells are permissive to neurite outgrowth and stabilize sensory axons during macrophage-induced axonal dieback after spinal cord injury. *J Neurosci*. 2010 Jan 6;30(1):255-65.
- Lane MA, Lee KZ, Salazar K, O'Steen BE, Bloom DC, Fuller DD, Reier PJ. Respiratory function following bilateral mid-cervical contusion injury in the adult rat. *Exp Neurol*. 2012 May;235(1):197-210.
- Valenzuela V, Collyer E, Armentano D, Parsons GB, Court FA, Hetz C. Activation of the unfolded protein response enhances motor recovery after spinal cord injury. *Cell Death Dis*. 2012 Feb 16;3:e272.
- McCreedy DA, Sakiyama-Elbert SE. Combination therapies in the CNS: engineering the environment.
- *Neurosci Lett*. 2012 Jun 25;519(2):115-21.
- 16. Johnson PJ, Tatara A, McCreedy DA, Shiu A, Sakiyama-Elbert SE. Tissue-engineered fibrin scaffolds containing neural progenitors enhance functional recovery in a subacute model of SCI. *Soft Matter*. 2010 Oct 21;6(20):5127-5137.
- Singh A, Balasubramanian S, Murray M, Lemay M, Houle J. Role of spared pathways in locomotor recovery after body-weight-supported treadmill training in contused rats. *J Neurotrauma*. 2011 Dec;28(12):2405-16.
- Singh A, Murray M, Houle JD. A training paradigm to enhance motor recovery in contused rats: effects of staircase training. *Neurorehabil Neural Repair*. 2011 Jan;25(1):24-34
- Luo J, Bo X, Wu D, Yeh J, Richardson PM, Zhang Y. Promoting survival, migration, and integration of transplanted Schwann cells by over-expressing polysialic acid. *Glia*. 2011 Mar;59(3):424-34.
- Wu D, Yang P, Zhang X, Luo J, Haque ME, Yeh J, Richardson PM, Zhang Y, Bo X. Targeting a dominant negative rho kinase to neurons promotes axonal outgrowth and partial functional recovery after rat rubrospinal tract lesion. *Mol Ther*. 2009 Dec;17(12):2020-30.

- Starkey ML, Bartus K, Barritt AW, Bradbury EJ. Chondroitinase ABC promotes compensatory sprouting of the intact corticospinal tract and recovery of forelimb function following unilateral pyramidotomy in adult mice. *Eur J Neurosci*. 2012
- James ND, Bartus K, Grist J, Bennett DL, McMahon SB, Bradbury EJ. Conduction failure following spinal cord injury: functional and anatomical changes from acute to chronic stages. *J Neurosci*. 2011 Dec 14;31(50):18543-55.
- Bartus K, James ND, Bosch KD, Bradbury EJ. Chondroitin sulphate proteoglycans: key modulators of spinal cord and brain plasticity. *Exp Neurol*. 2012 May;235(1):5-17.
- Thompson AK, Lapallo B, Duffield M, Abel BM, Pomerantz F. Repetitive common peroneal nerve stimulation increases ankle dorsiflexor motor evoked potentials in incomplete spinal cord lesions. *Exp Brain Res*. 2011
- Bosch KD, Bradbury EJ, Verhaagen J, Fawcett JW, McMahon SB. Chondroitinase ABC promotes plasticity of spinal reflexes following peripheral nerve injury. *Exp Neurol*. 2012 Nov;238(1):64-78
- Laskowski CJ, Bradke F. In vivo imaging: a dynamic imaging approach to study spinal cord regeneration. *Exp Neurol*. 2013 Apr;242:11-7.
- Hilton BJ, Assinck P, Duncan GJ, Lu D, Lo S, Tetzlaff W. Dorsolateral funiculus lesioning of the mouse cervical spinal cord at C4 but not at C6 results in sustained forelimb motor deficits. *J Neurotrauma*. 2013 Mar 21. [Epub ahead of print]
- Beller JA, Kulengowski B, Kobraei EM, Curinga G, Calulot CM, Bahrami A, Hering TM, Snow DM. Comparison of sensory neuron growth cone and filopodial responses to structurally diverse aggregan variants, in vitro. *Exp Neurol*. 2013 Mar 1;247C:143-157.
- McCall J, Weidner N, Blesch A. Neurotrophic factors in combinatorial approaches for spinal cord regeneration. *Cell Tissue Res*. 2012 Jul;349(1):27-37. doi: 10.1007/s00441-012-1388-6
- Dougherty BJ, Lee KZ, Gonzalez-Rothi EJ, Lane MA, Reier PJ, Fuller DD. Recovery of inspiratory intercostal muscle activity following high cervical hemisection. *Respir Physiol Neurobiol*. 2012 Sep 30;183(3):186-92. doi: 10.1016/j.resp.2012.06.006.
- Blackmore MG, Wang Z, Lerch JK, Motti D, Zhang YP, Shields CB, Lee JK, Goldberg JL, Lemmon VP, Bixby JL. Krüppel-like Factor 7 engineered for transcriptional activation promotes axon regeneration in the adult corticospinal tract. *Proc Natl Acad Sci U S A*. 2012 May 8;109(19):7517-22
- Kundi S, Bicknell R, Ahmed Z. The role of angiogenic and wound-healing factors after spinal cord injury in mammals. *Neurosci Res*. 2013 Apr 3. doi:pii: S0168-0102(13)00092-8.